

TOROIDAL AND BOUNDARY-REDUCING DEHN FILLINGS

C. MCA. GORDON AND J. LUECKE

ABSTRACT. Let M be a simple 3-manifold with a toral boundary component $\partial_0 M$. If Dehn filling M along $\partial_0 M$ one way produces a toroidal manifold and Dehn filling M along $\partial_0 M$ another way produces a boundary-reducible manifold, then we show that the absolute value of the intersection number on $\partial_0 M$ of the two filling slopes is at most two. In the special case that the boundary-reducing filling is actually a solid torus and the intersection number between the filling slopes is two, more is said to describe the toroidal filling.

1. INTRODUCTION

Following [W2], let us call a compact, orientable 3-manifold *simple* if it contains no essential sphere, disk, torus or annulus. Let M be such a manifold, and let $\partial_0 M$ be a torus component of ∂M . If γ is a *slope* on $\partial_0 M$ (the isotopy class of an essential unoriented circle), then as usual $M(\gamma)$ will denote the manifold obtained by γ -Dehn filling on M . Thus $M(\gamma) = M \cup V_\gamma$, where V_γ is a solid torus, glued to M by a homeomorphism from $\partial_0 M$ to ∂V_γ taking γ to the boundary of a meridian disk of V_γ .

If γ, δ are two slopes on $\partial_0 M$ such that $M(\gamma)$ and $M(\delta)$ are not simple, then there are several results giving upper bounds for $\Delta(\gamma, \delta)$ (the minimal geometric intersection number of γ and δ) for the various possible pairs of essential surfaces that arise, which in many cases are best possible (see [W2] for more details). In the present paper we dispose of one of the remaining cases, namely, that in which the surfaces in question are a torus and a disk.

Theorem 1.1. *Let M be a simple 3-manifold such that $M(\gamma)$ is toroidal and $M(\delta)$ is boundary-reducible. Then $\Delta(\gamma, \delta) \leq 2$.*

Examples showing that this bound is best possible are given in [HM1] and [MM].

It was previously known that $\Delta(\gamma, \delta) \leq 3$, by [W1]. We shall therefore assume from now on that $\Delta(\gamma, \delta) = 3$, and eventually obtain a contradiction. We shall also assume that ∂M has exactly two components $\partial_0 M$ and $\partial_1 M$, each a torus, and that M is a $\mathbb{Q}$ -homology cobordism between them, since otherwise $\Delta(\gamma, \delta) \leq 1$ by [W2, Theorem 4.1].

In [GLu3] it was shown that if $M(\gamma)$ is toroidal and $M(\delta) \cong S^3$ then $\Delta(\gamma, \delta) \leq 2$, and much of the proof of Theorem 1.1 consists of carrying over the arguments there to the

1991 *Mathematics Subject Classification.* Primary 57N10; Secondary 57M25.

Key words and phrases. Dehn surgery, Dehn filling, hyperbolic Dehn surgery.

present context. (In [GLu3] we analyzed the intersection of the punctured surfaces in M coming from an essential torus in $M(\gamma)$ and a Heegaard sphere in $M(\delta)$; here the Heegaard sphere is replaced by an essential disk.) This quickly leads to a proof of Theorem 1.1 when t , the number of points of intersection of the essential torus in $M(\gamma)$ with the core of V_γ , is at least 4. This is completed in Section 4. Sections 2, 3 and 4 parallel the corresponding Sections of [GLu3]; (in Section 2 it is also pointed out that Sections 6 and 7 of [GLu3] carry over without change). The main divergence in the proofs is in the case $t = 2$. In [GLu3] this case was handled by showing that the associated knot K in S^3 was strongly invertible, and then appealing to a result of Eudave-Muñoz [E-M]. Since no analog of this result is available in our present setting, we have to argue this case directly. This is done in Section 5.

In Section 6 we specialize to the case that the boundary-reducible filling of M is a solid torus. Infinitely many examples of simple 3-manifolds M and slopes γ, δ with $\Delta(\gamma, \delta) = 2$, $M(\gamma)$ toroidal, and $M(\delta)$ a solid torus, are given in [MM]. In these examples, there is a twice-punctured essential torus in M whose boundary slope is γ . Section 6 is devoted to showing this always happens:

Theorem 6.1. *Let M be a simple 3-manifold such that $M(\gamma)$ is toroidal, $M(\delta)$ is a solid torus, and $\Delta(\gamma, \delta) = 2$. Let K_γ be the core of the attached solid torus in $M(\gamma)$ and $\hat{T}$ be an essential torus in $M(\gamma)$ that intersects K_γ minimally. Then $|K_\gamma \cap \hat{T}| = 2$.*

We thank the referee for his helpful comments, and in particular for suggesting a simpler proof of Lemma 5.9.

2. THE GRAPHS OF INTERSECTION

Let $K_\gamma(K_\delta)$ be the core of V_γ (resp. V_δ). Let $\hat{T}$ be an essential torus in $M(\gamma)$. We assume that $\hat{T}$ meets K_γ transversely, and that $t = |\hat{T} \cap K_\gamma|$ is minimal (over all essential tori $\hat{T}$ in $M(\gamma)$). We may also assume that $\hat{T} \cap V_\gamma$ consists of meridian disks, so $T = \hat{T} \cap M$ is a punctured torus, with t boundary components, each having slope γ on $\partial_0 M$.

Similarly, let $\hat{Q}$ be an essential disk in $M(\delta)$, meeting K_δ transversely, with $q = |\hat{Q} \cap K_\delta|$ minimal over all essential disks in $M(\delta)$. Then $Q = \hat{Q} \cap M$ is a punctured disk, with one boundary component, $\partial\hat{Q}$, on $\partial_1 M$, and q boundary components on $\partial_0 M$, each with slope δ .

By standard arguments we may assume that:

- (i) Q meets T transversely, in properly embedded arcs and circles;
- (ii) each component of $\partial Q \cap \partial_0 M$ meets each component of ∂T in $\Delta = \Delta(\gamma, \delta)$ points;
- (iii) no arc component of $Q \cap T$ is boundary parallel in either Q or T ;
- (iv) no circle component of $Q \cap T$ bounds a disk in either Q or T .

Then, as described in [GLu3, Section 2], the arc components of $Q \cap T$ define graphs G_Q in $\hat{Q}$ and G_T in $\hat{T}$.

The definitions and terminology of [GLu1, Section 2] and [GLu3, Section 2], in particular the notion of a **q-type**, carry over to our present context. We assume familiarity with this terminology.

Note that since $\hat{Q}$ does not separate $M(\delta)$, the signs of the labels around the vertices of G_T (equivalently, the signs of the intersections of K_δ with $\hat{Q}$) do not necessarily alternate. Thus we may designate the corners of faces of G_T as either $++$, $--$, or $+-$, according to the signs of the corresponding pair of labels. However, we have the following lemma.

Lemma 2.1. *Let $\mathcal{D}$ be a set of disk faces of G_T representing all $\mathbf{q}$ -types. Then there exists $\mathcal{D}' \subset \mathcal{D}$ such that $\mathcal{D}'$ represents all $\mathbf{q}$ -types and all the corners appearing in faces in $\mathcal{D}'$ are $+-$.*

Proof. Let $\mathcal{C}_0$ be the set of $++$ and $--$ corners, and $\mathcal{C}_1$ the set of $+-$ corners, that appear in faces belonging to $\mathcal{D}$. Let τ_0 be the $\mathcal{C}_0$ -type defined by $\tau_0|(+ + \text{ corner}) = +$, $\tau_0|(- - \text{ corner}) = -$.

Now let τ_1 be any $\mathcal{C}_1$ -type, and let τ be the $(\mathcal{C}_0 \cup \mathcal{C}_1)$ -type (τ_0, τ_1) . By hypothesis, there exists a face $D \in \mathcal{D}$ which represents τ . We may assume, by definition of τ_0 , that the character (see [GLu1, p.386]) of each edge endpoint at a corner of D belonging to $\mathcal{C}_0$ is $+$. Since edges join points of opposite character, and since the edge endpoints at a corner in $\mathcal{C}_1$ have distinct characters, it follows that no corner of D can belong to $\mathcal{C}_0$. We thus obtain our desired subcollection $\mathcal{D}' \subset \mathcal{D}$ such that $\mathcal{D}'$ represents all $\mathcal{C}_1$ -types, and hence all $\mathbf{q}$ -types. $\square$

Theorem 2.2. *G_T does not represent all $\mathbf{q}$ -types.*

Proof. Let $\mathcal{D}$ be a set of disk faces of G_T representing all $\mathbf{q}$ -types. By Lemma 2.1, we can assume that all corners appearing in faces in $\mathcal{D}$ are $+-$, i.e., correspond to points of intersection of K_δ with $\hat{Q}$ of opposite sign. By [GLu1, Lemma 3.1], there exists $\mathcal{D}_0 \subset \mathcal{D}$ representing all $\mathbf{q}$ -types such that each face in $\mathcal{D}_0$ is locally on the same side of $\hat{Q}$. By [GLu2, Lemma 4.4], there exists $\mathcal{D}_1 \subset \mathcal{D}_0$ such that $\{[\partial D] : D \in \mathcal{D}_1\}$ is a basis for $\mathbb{R}^{c(\mathcal{D}_1)}$. Here $c(\mathcal{D}_1)$ is the number of corners appearing in faces belonging to $\mathcal{D}_1$, and $[\partial D]$ is the element of $\mathbb{Z}^{c(\mathcal{D}_1)} \subset \mathbb{R}^{c(\mathcal{D}_1)}$ (defined up to sign) obtained in the obvious way, by taking the algebraic sum of the corners in D . Tubing $\hat{Q}$ along the annuli in ∂V_δ corresponding to the corners appearing in faces belonging to $\mathcal{D}_1$ and surgering by the disks in $\mathcal{D}_1$ then gives a disk $\hat{Q}' \subset M(\delta)$ with $\partial \hat{Q}' = \partial \hat{Q}$, and $|\hat{Q}' \cap K_\delta| < |\hat{Q} \cap K_\delta| = q$, contradicting the minimality of q . $\square$

The definition of a *web* Λ in G_Q is exactly as in [GLu3, p.601]. If U is the component of $\hat{Q} - \text{nhd}(\Lambda)$ containing $\partial \hat{Q}$ then we say $D_\Lambda = \hat{Q} - U$ is the *disk bounded by* Λ . A *great web* is a web Λ such that Λ contains all the edges of G_Q that lie in D_Λ .

Exactly as in [GLu3, proof of Theorem 2.5], Theorem 2.2 implies

Theorem 2.3. *G_Q contains a great web.*

Since $\Delta = 3$, the arguments in [GLu3, Sections 6 and 7] apply here without change to show

Theorem 2.4. *$M(\gamma)$ does not contain a Klein bottle.*

Since $M(\delta)$ is boundary-reducible and $\gamma \neq \delta$, $M(\gamma)$ is irreducible by [S].

Finally we note that since M is a $\mathbb{Q}$ -homology cobordism between $\partial_0 M$ and $\partial_1 M$, $M(\gamma)$ is a $\mathbb{Q}$ -homology $S^1 \times D^2$. Hence $\hat{T}$ separates $M(\gamma)$, into X and X' , say. It follows that the faces of G_Q may be shaded alternately black and white, with the black faces lying in X and the white faces lying in X' .

3. SCHARLEMANN CYCLES AND EXTENDED SCHARLEMANN CYCLES

In this section we follow [GLu3, Section 3] and show that the relevant statements there hold in our present setting. For ease of reference, here and in Section 4 we will give lemmas etc. the same numbers as the corresponding statements in [GLu3].

Lemma 3.1. *The edges of a Scharlemann cycle in G_Q cannot lie in a disk in $\hat{T}$.*

Proof. Otherwise, as in [GLu3, proof of Lemma 3.1], $M(\gamma)$ would have a lens space summand, and hence be reducible. $\square$

Theorem 3.2. *If $t \geq 4$ then G_Q does not contain an extended Scharlemann cycle.*

Proof. The proof of [GLu3, Claim 3.3] goes through verbatim in our present context.

The proof of [GLu3, Claim 3.4] remains valid here once we note in addition that ∂M_1 cannot be parallel to $\partial M(\gamma)$ ($= \partial_1 M$). For then $\hat{T}$ would lie in a collar of $\partial M(\gamma)$ in $M(\gamma)$, contradicting the essentiality of $\hat{T}$.

We now follow the remainder of the proof of [GLu3, Theorem 3.2] as closely as possible.

Let $M_2 = \overline{X - M_1}$. Then $X = M_1 \cup_B M_2$. Similarly, since $M(\gamma)$ is a $\mathbb{Q}$ -homology $S^1 \times D^2$, A separates X' , into M_3 and M_4 , say.

Let T_2, T_3, T_4 be the tori defined as $T_2, \hat{T}_3, \hat{T}_4$ in [GLu3, pp.609–610]. These satisfy $|T_i \cap K_\gamma| < |\hat{T} \cap K_\gamma|$, $i = 2, 3, 4$, and hence, by the minimality of t , T_i is either compressible, and therefore (since $M(\gamma)$ is irreducible) bounds a solid torus in $M(\gamma)$, or is peripheral, i.e., parallel to $\partial M(\gamma)$, $i = 2, 3, 4$. It follows that exactly one of M_2, M_3, M_4 is a collar of $\partial M(\gamma)$ in $M(\gamma)$, and the other two are solid tori.

Now let T_0 be the torus $A \cup (C_1 - A_1) \cup B$. Then T_0 separates $M(\gamma)$ into $Y = M_1 \cup_A M_3$ and $Y' = M_2 \cup_{C_2} M_4$. Also $|T_0 \cap K_\gamma| < t$. Hence again T_0 either bounds a solid torus or is peripheral.

If T_0 is peripheral, then $M(\gamma) \cong Y$ or Y' , the union of two solid tori along an annulus essential in the boundary of each. Since $M(\gamma)$ is irreducible, this implies that $M(\gamma)$ is a Seifert fiber space over the disk with at most two singular fibers, contradicting the fact that it contains an essential torus.

If Y , say, is a solid torus, then M_2 or M_4 (say M_2) is a collar of $\partial M(\gamma)$, and hence $M(\gamma) \cong Y \cup M_4$ is a union of two solid tori, which is a contradiction as before. We get a similar contradiction if Y' is a solid torus. $\square$

We remark that we will only use Theorem 3.2 for extended Scharlemann cycles of length 2; see Section 4. However, we have given the proof of the general statement, for possible future reference.

Since $M(\gamma)$ does not contain a Klein bottle by Theorem 2.4, the proof of [GLu3, Lemma 3.10] applies verbatim to give the same statement here:

Lemma 3.10. *Let σ_1 and σ_2 be Scharlemann cycles of length 2 in G_Q on distinct label-pairs. Then the loops on $\hat{T}$ formed by the edges of σ_1 and σ_2 respectively are not isotopic on $\hat{T}$.*

Similarly, the proof of [GLu3, Theorem 3.11] gives

Theorem 3.11. *At most three labels can be labels of Scharlemann cycles of length 2 in G_Q .*

4. THE CASE $t \geq 4$

We obtain a contradiction in this case exactly as in [GLu3, Section 4].

Let Λ be a great web in G_Q , as guaranteed by Theorem 2.3, and D_Λ be the disk that it bounds. For every label x of G_Q let Λ_x be the graph in D_Λ consisting of the vertices of Λ along with all x -edges in Λ .

Let V be the number of vertices of Λ .

Lemma 4.2. *Let x be a label of G_Q . If Λ_x has at least $3V - 4$ edges, then Λ_x contains a bigon.*

Proof. The proof of [GLu3, Lemma 4.2] applies here without change.

We take this opportunity to thank Chuichiro Hayashi and Kimihiko Motegi for pointing out that the case where the face f is a monogon was not explicitly addressed in [GLu3]. However, in that case Λ_x could have at most one ghost x -label, since otherwise Λ would have more than t ghost labels. Hence $E \geq 3V - 1$. Also, if Λ_x does not contain a bigon then $1 + 3(F - 1) \leq 2E$. Therefore $2 = V - E + F \leq (E + 1)/3 - E + 2(E + 1)/3 = 1$, a contradiction. $\square$

Lemma 4.2 and Theorems 3.2 and 3.11 then lead to a contradiction exactly as in [GLu3, p.615].

5. THE CASE $t = 2$

As in the previous section, let Λ be a great web in G_Q .

Lemma 5.1. *Λ contains a bigon.*

Proof. Regard Λ as a graph in the disk, with V vertices, E edges, and F faces.

Since Λ has at most two ghost labels, $2E \geq 6V - 2$, so $E \geq 3V - 1$. Suppose Λ contains no bigon; then $3F < 2E$. Therefore $1 = V - E + F < (E + 1)/3 - E + 2E/3 = \frac{1}{3}$, a contradiction. $\square$

By the parity rule, each edge of Λ must connect different vertices of G_T when considered as edges lying in G_T . Recall that there are four *edge classes* in G_T , i.e., isotopy classes of non-loop edges of G_T in $\hat{T} \text{ rel}\{\text{vertices of } G_T\}$, which we call $1, \alpha, \beta, \alpha\beta$, as illustrated in [GLu3, Figure 7.1]. The ordering of these classes around vertices 1 and 2 of G_T is indicated in Figure 5.1. We may label an edge e of G_Q by the class of the corresponding edge of G_T ; we refer to this label as the *edge class label* of e .




FIGURE 5.1

Lemma 5.2. *Any two black (white) bigons in Λ have the same pair of edge class labels.*

Proof. Let f, f' be two (say) black bigons in Λ , with vertices x, x' and y, y' , and edge class labels λ, μ and λ', μ' , respectively. See Figure 5.2. Note that $\lambda \neq \mu$ and $\lambda' \neq \mu'$ by Lemma 3.1.




FIGURE 5.2

Let H_{12} be the 1-handle $V_\gamma \cap X$. The anticlockwise ordering of the labels x, x', y, y' around vertex 1 of G_T must agree with their clockwise ordering around vertex 2, since the corners of f and f' give four disjoint arcs on the annulus $\partial H_{12} - \hat{T}$ joining corresponding labels. However, if $\lambda = \lambda'$ but $\mu \neq \mu'$ then it is easy to see that these orderings do not agree. For example, the case $\lambda = \lambda' = 1$, $\mu = \alpha$, $\mu' = \beta$ is illustrated in Figure 5.3. Similarly, the pairs $\{1, \alpha\beta\}$, $\{\alpha, \beta\}$ give incompatible orderings of the labels.




FIGURE 5.3

Thus the only possible distinct pairs of edge class labels for f and f' are $\{1, \alpha\}$ and $\{\alpha\beta, \beta\}$ (or $\{1, \beta\}$ and $\{\alpha\beta, \alpha\}$). Now by shrinking H_{12} to its core, $H_{12} \cup f$ becomes a Möbius band B in X such that ∂B is the loop on $\hat{T}$ formed by the edges of f . Similarly f' gives rise to a Möbius band B' . If the edge class labels for f and f' are as above, then ∂B and $\partial B'$ may be isotoped on $\hat{T}$ to be disjoint. This produces a Klein bottle in $M(\gamma)$, contradicting Theorem 2.4. $\square$

Lemma 5.3. *Let e, e' be edges of Λ incident to a vertex v of Λ , with the same label at v . Then e and e' have distinct edge class labels.*

Proof. If not, then e and e' would be parallel in G_T , and hence would cobound a family of

$q + 1$ parallel edges of G_T . The argument of [GLi, p.130, Case (2)] now constructs a cable space in M , contradicting the assumption that M is simple. $\square$

If f is a white (black) face of Λ such that each edge of f belongs to a black (resp. white) bigon, then we say that f is *surrounded by bigons*.

Lemma 5.4. *Λ does not contain a face surrounded by bigons.*

Proof. Suppose that Λ contains, say, a white face f surrounded by black bigons.

By Lemma 5.2 (and Lemma 3.1) all the black bigons have the same pair of (distinct) edge class labels λ, μ .

Let v be a vertex of f . By Lemma 5.3 the edges of the two black bigons incident to v having label 1 (say) at v have distinct edge class labels. Therefore the two edges of f incident to v have the same edge class label (see Figure 5.4). Hence all the edges of f have the same edge class label, contradicting Lemma 3.1. $\square$




FIGURE 5.4

Lemma 5.5. *Not all black faces of Λ are bigons. Similarly, not all white faces of Λ are bigons.*

Proof. Suppose, for example, that all black faces of Λ are bigons.

Define a graph Σ in the disk $\hat{Q}$ as follows. The vertices of Σ are the vertices of Λ , and the edges of Σ are in one-one correspondence with the black bigons of Λ , each edge joining the vertices at the corners of the corresponding bigon and lying in its interior. Then each vertex of Σ has valency at least 2, except that if Λ has a vertex at which there are two ghost labels, then that vertex may have valency 1 in Σ . Hence if V and E are the number of vertices and edges of Σ respectively, then $2E \geq 2V - 1$, and so $2E \geq 2V$. Therefore

$$\begin{aligned} 1 &\leq V - E + \sum \chi(f) \text{ (the sum being over all faces } f \text{ of } \Sigma) \\ &\leq \sum \chi(f), \end{aligned}$$

implying that Σ has a disk face. But such a face corresponds to a white face of Λ surrounded by black bigons, contradicting Lemma 5.4. $\square$

We shall say that two faces f_1, f_2 of Λ of the same color are *isomorphic* if the cyclic sequences of edge class labels obtained by reading around the boundaries of f_1 and f_2 in the same direction, are equal.

Lemma 5.6. *Either all black faces of Λ are isomorphic or all white faces of Λ are isomorphic.*

Proof. Recall that $\hat{T}$ separates $M(\gamma)$, into X and X' . We suppose without loss of generality that $\partial M(\gamma) \subset X$.

Recall also that H_{12} is the 1-handle $V_\gamma \cap X$. Let f be any black face of Λ , and let $N = \text{nhd}(\hat{T} \cup H_{12} \cup f) \subset X$. Then (since all the vertices of Λ have the same sign) $\partial N = \hat{T} \cup T_1$, where T_1 is a torus. Since $T_1 \cap K_\gamma = \emptyset$, and since M is irreducible and atoroidal, T_1 is parallel to $\partial_1 M = \partial M(\gamma)$. Hence $W = \overline{X - H_{12}}$ is a compression body, with $\partial_+ W$ the genus 2 surface obtained by adding the tube $\partial H_{12} - \hat{T}$ to T , and $\partial_- W = \partial M(\gamma)$. It follows that f is the unique non-separating disk in W , up to isotopy.

Let f' be any other black face of Λ . Then ∂f and $\partial f'$ are isotopic in $\partial_+ W$, and hence (freely) homotopic in $\hat{T} \cup H_{12}$. Now $\pi_1(\hat{T} \cup H_{12}) \cong \pi_1(\hat{T}) * \mathbb{Z}$, where, taking as base-“point” a disk neighborhood in $\hat{T}$ of an edge in G_T in edge class 1, $\pi_1(\hat{T}) \cong \mathbb{Z} \times \mathbb{Z}$ has basis $\{\alpha, \beta\}$, represented by edges in the correspondingly named edge classes, oriented from vertex 2 to vertex 1, and $\mathbb{Z}$ is generated by x , say, represented by an arc in H_{12} going from vertex 1 to vertex 2. Then, if the sequence of edge class labels around ∂f (in the appropriate direction) is $(\gamma_1, \dots, \gamma_n)$, ∂f represents $\gamma_1 x \gamma_2 x \dots \gamma_n x \in \pi_1(\hat{T}) * \mathbb{Z}$, and similarly for $\partial f'$. Since ∂f and $\partial f'$ are homotopic in $\hat{T} \cup H_{12}$, we conclude that the corresponding cyclic sequences $(\gamma_1, \dots, \gamma_n)$ and $(\gamma'_1, \dots, \gamma'_n)$ are equal, i.e., f and f' are isomorphic. Thus all black faces of Λ are isomorphic, and the lemma is proved. $\square$

By Lemma 5.1, Λ contains a bigon. We may therefore assume from now on that Λ contains a black bigon.

Lemma 5.7. *All white faces of Λ are isomorphic.*

Proof. If not, then by Lemma 5.6 all black faces of Λ are isomorphic, and therefore bigons. But this contradicts Lemma 5.5. $\square$

Lemma 5.8. *Λ does not contain a white bigon.*

Proof. This follows from Lemmas 5.7 and 5.5. $\square$

Recall that D_Λ is the disk bounded by the great web Λ . Travelling around ∂D_Λ in some direction defines in the obvious way a cyclic sequence of edges of Λ , $(e_1, \dots, e_n)$. (Note that the same edge may appear twice, with opposite orientations.) If the subgraph of Λ consisting of these edges has no cut vertex, define $\Lambda_0 = \Lambda$. (Recall that a cut vertex of a connected graph is one whose removal disconnects the graph.) Otherwise, let v_0 be an outermost cut vertex of this subgraph. This vertex cuts off a subdisk of D_Λ which is the disk D_{Λ_0} bounded by a subgraph Λ_0 of Λ (see Figure 5.5). We shall call Λ_0 an *extremal subgraph* of Λ . If $\Lambda_0 \neq \Lambda$ we call v_0 the *attaching vertex* of Λ_0 .




FIGURE 5.5

An *interior vertex* of Λ is a vertex of Λ that is not an endpoint of any of the edges $e_1, \dots, e_n$. All six corners at such a vertex belong to faces of Λ .

Lemma 5.9. *Λ contains an interior vertex.*

Proof. The following proof was suggested by the referee.

Since Λ has at most two ghost labels, Λ has an extremal subgraph Λ_0 such that either $\Lambda_0 = \Lambda$ or Λ_0 contains at most one ghost label of Λ at a vertex other than the attaching vertex v_0 of Λ_0 . If $\Lambda = \Lambda_0$, we can also choose a vertex v_0 of Λ such that Λ has at most one ghost label at a vertex other than v_0 . We shall show that Λ_0 contains an interior vertex.

Suppose for contradiction that Λ_0 has no interior vertices. Let m be the number of vertices of Λ_0 . Note that $m > 1$, since G_Q has no trivial loops. Let $\overline{\Lambda}_0$ be the reduced graph of Λ_0 , obtained by amalgamating each family of parallel edges to a single edge. If $m = 2$ or 3 , then each vertex of $\overline{\Lambda}_0$ has valency 1 or 2 respectively. If $m \geq 4$, the interior edges of $\overline{\Lambda}_0$ decompose D_{Λ_0} into subdisks, and by considering outermost such subdisks we see that $\overline{\Lambda}_0$ has at least two vertices of valency 2. Hence in all cases $\overline{\Lambda}_0$ has a vertex $v \neq v_0$ of valency 1 or 2. Since there is at most one ghost label at v , Λ_0 has a family of at least three parallel edges at v , and hence contains a white bigon, contradicting Lemma 5.8. $\square$

Lemma 5.10. *Each white face of Λ has at least three distinct edge class labels.*

Proof. By Lemma 5.9, Λ contains an interior vertex v . Then all three white corners at v belong to faces of Λ . By Lemma 5.3, the three edges with label (say) 1 at v have distinct edge class labels. The result now follows from Lemma 5.7. $\square$

Lemma 5.11. *Each white face of Λ has length at least 4.*

Proof. The argument in the first paragraph of [GLu3, proof of Lemma 3.7] shows that a face of Λ of length 3 has only two distinct edge class labels, contradicting Lemma 5.10. $\square$

Recall that Λ has either 0 or 2 ghost labels. In the former case, let $\Lambda^+ = \Lambda$. In the latter case, let $\Lambda^+ \subset \hat{Q}$ be obtained from Λ by adjoining two disjoint arcs running from the ghost labels to $\partial\hat{Q}$. The complementary regions of Λ^+ in $\hat{Q}$ are the faces of Λ together with either one or two *outside regions*, i.e., regions that meet $\partial\hat{Q}$. The black/white shading of the faces of Λ extends to a shading of the complementary regions of Λ^+ .

Define a graph Γ in the disk $\hat{Q}$ as follows. The vertices of Γ are the (*fat*) vertices of Λ , together with *dual vertices*, one in the interior of each black complementary region of Λ^+

that is either a face of Λ of length at least 3, or an outside region. The edges of Γ consist of edges joining each dual vertex to the fat vertices in the boundary of the corresponding region, together with an edge in the interior of each black bigon in Λ , joining the (fat) vertices at the ends of the bigon. (If $\Lambda^+ = \Lambda$ and the outside region is black, then there is a choice involved in defining the edges incident to the dual vertex corresponding to that region.)

Lemma 5.12. *Γ has a face of length at most 5.*

Proof. The valency of each fat vertex of Γ is 3. The valency of each dual vertex corresponding to a face of Λ is at least 3. There is at most one dual vertex corresponding to an outside region, and that vertex has valency at least 1. Hence, if Γ has V vertices and E edges then $2E \geq 3V - 2$. If Γ has F faces, and each face has length at least 6, then $6F < 2E$. Therefore $1 = V - E + F < 2(E + 1)/3 - E + E/3 = \frac{2}{3}$, a contradiction. $\square$

Let g be a face of Γ as in Lemma 5.12. Then g contains a unique white face f of Λ . If f has b edges which are edges belonging to black bigons of Λ , and c other edges, then the length of g is $b + 2c \leq 5$. On the other hand $b + c \geq 4$ by Lemma 5.11. Also, $c > 0$ by Lemma 5.4; hence $c = 1$. Thus every edge of f except one belongs to a black bigon, and the argument in the proof of Lemma 5.4 shows that every such edge has the same edge class label. Therefore f has at most two edge class labels, contradicting Lemma 5.10.

We have thus shown that the case $t = 2$ is impossible.

This completes the proof of Theorem 1.1.

6. KNOTS IN SOLID TORI

In [GLu3] and [GLu4] it is shown that if M is simple, $M(\gamma)$ is toroidal, $M(\delta)$ is S^3 , and $\Delta(\gamma, \delta) = 2$, then $t = 2$. [MM] gives examples of simple manifolds, M , for which $M(\gamma)$ is toroidal, $M(\delta) = S^1 \times D^2$, and $\Delta(\gamma, \delta) = 2$. In these examples $t = 2$. It is natural then to ask if this is always the case.

Theorem 6.1. *Let M be simple 3-manifold such that $M(\gamma)$ is toroidal, $M(\delta)$ is $S^1 \times D^2$, and $\Delta(\gamma, \delta) = 2$. Let K_γ be the core of the attached solid torus in $M(\gamma)$ and $\hat{T}$ be an essential torus in $M(\gamma)$ that intersects K_γ minimally. Then $|K_\gamma \cap \hat{T}| = 2$.*

The remainder of this section is devoted to the proof of Theorem 6.1. Let $\lambda = S^1 \times *$, $\mu = * \times \partial D^2$ in $\partial_1 M = \partial M(\delta)$ be a longitude and meridian respectively of $S^1 \times D^2$.

For any integer n , let M_n be the $(\lambda + n\mu)$ -filling of M .

Lemma 6.2. *M_n is simple for n sufficiently large.*

Proof. By [Th], M is hyperbolic and all but finitely many fillings on a single component of ∂M are hyperbolic. See also Theorem 1.3 of [G]. $\square$

Let $N = M(\gamma)$, and parametrize slopes (as rational numbers) on $\partial N = \partial_1 M$ using λ, μ . Thus $M_n(\gamma) = N(n)$.

Let $\hat{T}$ be an essential torus in N that minimizes $|K_\gamma \cap \hat{T}|$.

Lemma 6.3. *If $\hat{T}$ compresses in $N(n_1)$ and $N(n_2)$ then either $|K_\gamma \cap \hat{T}| = 2$ or $|n_1 - n_2| \leq 2$.*

Proof. We assume for contradiction that every essential torus in N (which must separate since $\Delta(\gamma, \delta) = 2$) intersects K_γ at least four times and that $\hat{T}$ compresses in $N(n_1), N(n_2)$ where $|n_1 - n_2| \geq 3$. Applying [CGLS, Theorem 2.4.2] to the side of $\hat{T}$ containing $\partial_1 M$, this last condition guarantees an essential annulus $A' \subset N$ such that one boundary component of A' is $A' \cap \hat{T}$, the other is $A' \cap \partial_1 M$, and $A' \cap \partial_1 M$ is distance 1 from each of the integral slopes n_1 and n_2 . Since $|n_1 - n_2| \geq 3$, $A' \cap \partial_1 M$ must be a copy of μ . Surgering $\hat{T}$ along A' gives a properly embedded annulus $A'' \subset N$ whose boundary consists of two copies of μ . Since $\hat{T}$ is essential in N , so is A'' .

Let A be an essential annulus in N , with boundary consisting of two copies of μ , such that $|K_\gamma \cap A|$ is minimal over all such annuli. Since M contains no essential annulus, $K_\gamma \cap A \neq \emptyset$. Let D be a meridian disk of $M(\delta) = S^1 \times D^2$, chosen so as to minimize $|K_\delta \cap D|$. We may isotope D so that $\partial D \cap \partial A = \emptyset$. Then $T' = A \cap M$ and $Q = D \cap M$ are essential planar surfaces in M , whose “outer” boundaries, ∂A and ∂D lie on $\partial_1 M$ and are disjoint, and whose “inner” boundaries $\partial T' \cap \partial_0 M$, $\partial Q \cap \partial_0 M$ consist of, say, t' copies of γ and q copies of δ respectively.

Recall that $\Delta(\gamma, \delta) = 2$. As in Section 2 the arc components of $T' \cap Q$ give rise to graphs $G_{T'}, G_Q$ in A, D (resp.). By abstractly identifying the two boundary components of A , we may regard $G_{T'}$ as a graph in a torus, and we are now exactly in the combinatorial set-up of Section 2. In particular we may apply Theorem 2.3 to conclude that G_Q contains a great web, Λ . (Note that the one face of $G_{T'}$ in the torus which is not a face of $G_{T'}$ in A is not a disk face, hence would not be involved in any collection of faces representing all types). Because Λ has at most t' ghost labels and $\Delta(\gamma, \delta) = 2$, there is a label x of G_Q and a vertex y_0 of Λ such that the following conditions are satisfied:

- (i) for any vertex y of Λ other than y_0 , there is an edge of Λ incident to y at each occurrence of the label x on y , and
- (ii) there is an edge of Λ incident to y_0 at some occurrence of the label x at y_0 .

If we let Σ be the set of all x -edges of Λ , then Σ is a *great x -web* in the sense of [GLu2, p.390]. The argument of Theorem 2.3 of [GLu2] now shows that A is separating and either (a) G_Q must contain Scharlemann cycles on distinct label pairs or (b) $t' = 2$. (This argument uses Lemma 2.2 of the same paper, which requires the fact that consecutive labels on G_Q represent vertices of $G_{T'}$ of opposite sign. This assumption was shown not to be necessary in [HM2, Proposition 5.1]. However, one may guarantee this condition in the present context by showing that A is separating. This is done in the same way as in Theorem 2.3: show that Λ contains a Scharlemann cycle, then use the face of Λ bounded by this Scharlemann cycle to tube and compress A to get a new annulus A' such that $|A' \cap K_\delta| < |A \cap K_\delta|$. If A were non-separating, A' would be also, contradicting the minimality of A .)

Lemma 3.1 of [HM1] shows that conclusion (a) contradicts the minimality of $|A \cap K_\gamma|$. Thus we must have that $t' = 2$. Let X_1 and X_2 be the components into which A separates N . Then each ∂X_i is a torus meeting K_γ twice, and hence is compressible. Since N is

irreducible by [S], X_i must be a solid torus. As $N = X_1 \cup_A X_2$, N is Seifert-fibered over the disk with at most two exceptional fibers. But this contradicts the fact that N contains an essential torus. This contradiction finishes the proof of Lemma 6.3. $\square$

Remark. In fact, the possibility that N contains an essential torus intersecting K_γ twice can be excluded from the conclusion of Lemma 6.3. For example one can argue as above to show the existence of A , then argue as in [HS] to arrive at a contradiction. In particular, Sections 4 and 5 of that paper along with Lemma 3.1 of [HM1] show that the great x -web Λ that we see in the above argument cannot occur.

Proof of Theorem 6.1. Lemmas 6.2 and 6.3 show that if $|K_\gamma \cap \hat{T}| \neq 2$, then there is an integer J such that for all $n > J$, M_n is a simple manifold and $M_n(\gamma)$ contains an essential torus. On the other hand $M_n(\delta) = S^3$, so M_n is the exterior of a simple knot in S^3 . By Theorem 1.2 of [GLu3], (which is proved in [GLu3] and [GLu4]), each M_n contains an essential, twice-punctured torus, T_n , where ∂T_n is two copies of γ . If, for some n , T_n can be isotoped to lie inside M , then we are done. So assume not. Then for each $n > J$, there is an incompressible, ∂ -incompressible surface T'_n in M whose boundary is a non-empty collection of copies of $\lambda + n\mu$ on $\partial_1 M$ and two copies of γ on $\partial_0 M$. But this contradicts [H]. $\square$

In the above proof of Theorem 6.1, the direct appeal to Theorem 1.2 of [GLu3] was possible because of the assumption that $M(\delta)$ is a solid torus. Conceivably Theorem 6.1 is still true if it is only assumed that $M(\delta)$ is boundary-reducible. If so, a proof in this generality might be obtainable by suitably modifying the proof, in [GLu3] and [GLu4], of Theorem 1.2 of [GLu3].

REFERENCES

- [E-M] M. Eudave-Muñoz, *Essential tori obtained by surgery on a knot*, Pacific J. Math. **167** (1995), 81–117.
- [G] C. McA. Gordon, *Boundary slopes of punctured tori in 3-manifolds*, Trans. Amer. Math. Soc. (to appear).
- [GLi] C. McA. Gordon and R.A. Litherland, *Incompressible planar surfaces in 3-manifolds*, Topology and its Applications **18** (1984), 121–144.
- [GLu1] C. McA. Gordon and J. Luecke, *Knots are determined by their complements*, J. Amer. Math. Soc. **2** (1989), 371–415.
- [GLu2] C. McA. Gordon and J. Luecke, *Reducible manifolds and Dehn surgery*, Topology **35** (1996), 385–409.
- [GLu3] C. McA. Gordon and J. Luecke, *Dehn surgeries on knots creating essential tori, I*, Communications in Analysis and Geometry **3** (1995), 597–644.
- [GLu4] C. McA. Gordon and J. Luecke, *Dehn surgeries on knots creating essential tori, II*, Communications in Analysis and Geometry (to appear).
- [H] A.E. Hatcher, *On the boundary curves of incompressible surfaces*, Pacific J. Math. **99** (1982), 373–377.
- [HM1] C. Hayashi and K. Motegi, *Dehn surgery on knots in solid tori creating essential annuli*, Trans. Amer. Math. Soc. (to appear).
- [HM2] C. Hayashi and K. Motegi, *Only single twist on unknots can produce composite knots*, Trans. Amer. Math. Soc. (to appear).
- [HS] C. Hayashi and K. Shimokawa, *Symmetric knots satisfy the cabling conjecture*, preprint.

- [MM] K. Miyazaki and K. Motegi, *Toroidal and annular Dehn surgeries of solid tori*, preprint.
- [S] M. Scharlemann, *Producing reducible 3-manifolds by surgery on a knot*, *Topology* **29** (1990), 481–500.
- [Th] W. Thurston, *Three dimensional manifolds, Kleinian groups and hyperbolic geometry*, *Bull. Amer. Math. Soc.* **6** (1982), 357–381.
- [W1] Y-Q. Wu, *Dehn surgeries producing reducible manifold and toroidal manifold*, *Topology* (to appear).
- [W2] Y-Q. Wu, *Sutured manifold hierarchies, essential laminations, and Dehn surgery*, preprint.

DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF TEXAS AT AUSTIN, AUSTIN, TX 78712-1082
E-mail address: `gordon@math.utexas.edu`; `luecke@math.utexas.edu`